

# DcR1 / TRAILR3 Antibody, Rabbit PAb, Antigen Affinity Purified



Sino Biological  
Biological Solution Specialist

Catalog Number: 310555-T38

## GENERAL INFORMATION

<b>Immunogen:</b>	E. coli-derived Human DcR1 / TRAILR3 fragment
<b>Preparation</b>	Produced in rabbits immunized with E. coli-derived Human DcR1 / TRAILR3 fragment, and purified by antigen affinity chromatography.
<b>Ig Type:</b>	Rabbit IgG
<b>Specificity:</b>	Human DcR1 / TRAILR3
<b>Formulation:</b>	PBS, pH7.0 with 0.03% Proclin300
<b>Storage:</b>	This antibody can be stored at 2°C-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Avoid repeated freeze-thaw cycles.
<b>Alternative Names:</b>	TNFRSF10C

## APPLICATIONS

<b>Applications:</b>	WB, ICC/IF, IP
----------------------	----------------

## RECOMMENDED CONCENTRATION

<b>ICC/IF</b>	ICC/IF: 1:100-1:500
<b>Western Blot</b>	WB: 1:500-1:2000
<b>Immunoprecipitation</b>	IP:1-5μL/mg of lysate

*Please Note: Optimal concentrations/dilutions should be determined by the end user.*

**For Research Use Only. Not for use in diagnostic or therapeutic procedures.**

**Tel: +86-400-890-9989 (Global), +1-215-583-7898 (USA), +49(0)6196 9678656 (Europe)**


**Website: <http://www.sinobiological.com>**

# DcR1 / TRAILR3 Antibody, Rabbit PAb, Antigen Affinity Purified




Sino Biological  
Biological Solution Specialist

Catalog Number: 310555-T38



Immunofluorescence staining of TRAILR3 in A549 cells. Cells were fixed with 4% PFA, permeabilized with 0.3% Triton X-100 in PBS, blocked with 10% serum, and incubated with rabbit anti-Human TRAILR3 polyclonal antibody (dilution ratio 1:200) at 4°C overnight. Then cells were stained with the Alexa Fluor®488-conjugated Goat Anti-rabbit IgG secondary antibody (green) and counterstained with DAPI (blue). Positive staining was localized to Cytoplasm.




Anti-DcR1/TRAILR3 rabbit polyclonal antibody at 1:500 dilution  
Lane A: Hela Whole Cell Lysate  
Lane B: Jurkat Whole Cell Lysate  
Lane C: A549 Whole Cell Lysate

Lysates/proteins at 30 µg per lane.  
Secondary  
Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution.

Developed using the ECL technique.  
Performed under reducing conditions.

Predicted band size: 65 kDa  
Observed band size: 70 kDa



TNFRSF10C was immunoprecipitated using:  
Lane A: 0.5 mg Hela Whole Cell Lysate

4 µL anti-TNFRSF10C rabbit polyclonal antibody and 60 µg of Immunomagnetic beads Protein A/G.

Primary antibody:  
Anti-TNFRSF10C rabbit polyclonal antibody, at 1:100 dilution

Secondary antibody:  
Goat Anti-Rabbit IgG (H+L)/HRP at 1/10000 dilution

Developed using the ECL technique.  
Performed under reducing conditions.

Predicted band size: 65 kDa  
Observed band size: 65 kDa

For Research Use Only. Not for use in diagnostic or therapeutic procedures.

Tel: +86-400-890-9989 (Global), +1-215-583-7898 (USA), +49(0)6196 9678656 (Europe)

Website: <http://www.sinobiological.com>