

# Influenza B (B/Brisbane/60/2008) Neuraminidase / NA (Active)

Catalog Number: 40203-VNAHC



Sino Biological  
Biological Solution Specialist

## General Information

### Gene Name Synonym:

Neuraminidase, NA

### Protein Construction:

A DNA sequence encoding Influenza B (B/Brisbane/60/2008) Neuraminidase (4CPL\_A) (Met1-Leu466), termed as NA, was expressed.

**Source:** Influenza B

**Expression Host:** HEK293 Cells

## QC Testing

### Bio-Activity:

Measured by its ability to cleave a fluorogenic substrate, 2'-(4-Methylumbelliferyl)- $\alpha$ -D-N-acetylneuraminic acid. The specific activity is > 100 U. The specific activity is > 1000 U.

### Endotoxin:

< 1.0 EU per  $\mu$ g of the protein as determined by the LAL method

### Stability:

Samples are stable for up to twelve months from date of receipt at -70 °C

### Molecular Mass:

The recombinant influenza B (B/Brisbane/60/2008) Neuraminidase consists of 466 amino acids and predicts a molecular mass of 51.1 kDa.

### Formulation:

Lyophilized from sterile PBS, 1% Triton X-100 PH 7.4

Normally 5 % - 8 % trehalose and mannitol are added as protectants before lyophilization. Specific concentrations are included in the hardcopy of COA. Please contact us for any concerns or special requirements.

## Usage Guide

### Storage:

Store it under sterile conditions at -20°C to -80°C upon receiving. Recommend to aliquot the protein into smaller quantities for optimal storage.

### Avoid repeated freeze-thaw cycles.

### Reconstitution:

It is recommended that 1 ml sterile water be added to the vial to prepare a stock solution.

## Protein Description

Neuraminidase (NA) is a major membrane glycoproteins found on the surface of influenza virus. NA specifically catalyzes the hydrolysis removal of terminal sialic acid residues from viral and cellular glycoconjugates. It is known that HA binds to the sialic acid-containing receptors on the surface of host cells during initial infection, and at the end of an infectious cycle, NA cleaves the HA-sialic acid bondage from the newly formed virions and the host cell receptors during budding. NA thus is described as a receptor-destroying enzyme which facilitates virus release and efficient spread of the progeny virus from cell to cell. NA is a single-pass type I I membrane protein which exists as a homotetramer, and the transmembrane domain is involved in lipid raft association during intracellular transport. NA is suggested to play a role in the determination of host range restriction on replication and virulence. Nine subtypes of NA have been identified, and subtypes N1 and N2 have been positively linked to epidemics in man. Influenza A H1N1 virus is a subtype of influenza A virus. Some strains of H1N1 are endemic in humans and cause a small fraction of all influenza-like illness and a small fraction of all seasonal influenza. H1N1 strains caused a few percent of all human flu infections in 2004-2005. Other strains of H1N1 are endemic in pigs (swine influenza) and in birds (avian influenza). H1N1 was the most common cause of human influenza (flu) in 2009. In June 2009, the World Health Organization declared the new strain of swine-origin H1N1 as a pandemic. This strain is often called swine flu by the public media. This novel virus spread worldwide and had caused about 17,000 deaths by the start of 2010.

## References

1. Barman, S. et al., 2000, J. Virol. 74: 6538-6545.
2. Colman, PM. et al., 1983, Nature. 303: 41-44.
3. Suzuki, T. et al., 2005, J. Virol. 79: 11705-11715.
4. von, Itzstein, M. 2007, Nat. Rev. Drug. Discov. 6: 967-974.

For Research Use Only. Not for use in diagnostic or therapeutic procedures.

Tel: +86-400-890-9989 (Global), +1-215-583-7898 (USA), +49(0)6196 9678656 (Europe)

Website: <http://www.sinobiological.com>