

Rat Tie2 / TEK HEK293 Cell Lysate (WB positive control)



Sino Biological Inc.
Biological Solution Specialist

Catalog Number: 80063-R02HL

TIE2 Transfected / Overexpression Cell Lysate Product Information

| | |
|------------------------------|--|
| Expressed Host: | HEK293 Cells |
| Products Description: | Human Cell lysate that Rat Tie2 / TEK transfected / overexpressed for Western blot (WB) positive control. The whole cell lysate is provided in 1X Sample Buffer (1X modified RIPA buffer+1X SDS loading buffer). |
| Sequence information: | A DNA sequence encoding the rat TEK (NP_001099207.1) extracellular domain (Met 4-Leu 743) was fused with the Fc region of human IgG1 at the C-terminus. |
| Predicted N Terminal: | Ala 23 |
| Molecule Mass: | The recombinant rat TEK/Fc is a disulfide-linked homodimer. The reduced monomer comprises 962 amino acids and predicts a molecular mass of 108 kDa. The apparent molecular mass of the rat TEK/Fc monomer is approximately 120-130 kDa in SDS-PAGE under reducing conditions due to glycosylation. |
| Species: | Rat |

TIE2 Transfected / Overexpression Cell Lysate Usage Guide

| | |
|---------------------------------|--|
| Preparation Method: | Cell lysate was prepared by homogenization in ice-cold modified RIPA Lysis Buffer with cocktail of protease inhibitors (Sigma). Cell debris was removed by centrifugation. Protein concentration was determined by Bradford assay (Bio-Rad protein assay, Microplate Standard assay). The cell lysate was boiled for 5 min in 1 x SDS loading buffer (50 mM Tris-HCl pH 6.8, 12.5% glycerol, 1% sodium dodecylsulfate, 0.01% bromophenol blue) containing 5% b-mercaptoethanol, and lyophilized. |
| Lysis Buffer: | Modified RIPA Lysis Buffer: 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 1mM EDTA, 1% Triton X-100, 0.1% SDS, 1% Sodium deoxycholate, 1mM PMSF. |
| Quality Control Testing: | 12.5% SDS-PAGE Stained with Coomassie Blue after protein purification. |
| Stability: | Samples are stable for up to twelve months from date of receipt. |
| Recommend Usage: | <ol style="list-style-type: none">1. Centrifuge the tube for a few seconds and ensure the pellet at the bottom of the tube.2. Re-dissolve the pellet using 200μL pure water and boil for 2-5 min.3. Store the lyophilized cell lysate at 4°C. After re-dissolution, recommend to aliquot it into smaller quantities and store at -80°C. |
| Storage Buffer: | 1 X Sample Buffer (1 X modified RIPA buffer+1 X SDS loading buffer). |
| Storage Instruction: | Store at 4°C. After re-dissolution, aliquot and store at -80°C. |
| Application notes: | Western blot (WB): Use at an assay dependent dilution. Other Applications: Not tested. Optimal dilutions/concentrations should be determined by the end user. |

Manufactured By Sino Biological Inc., FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

For US Customer: Fax: 267-657-0217

● Tel: 215-583-7898

Global Customer: Fax :+86-10-5862-8288

● Tel:+86-400-890-9989 ●

● <http://www.sinobiological.com>